

Forest structure characteristics on soil carbon and nitrogen storage of *Pinus massoniana* plantations in southern subtropic region

Kexin Zhang^{1,2}, Dongli Gao^{1,4}, Hong Guo¹, Ji Zeng³ and Xianzhao Liu^{1,2*}

(1 Research Institute of Forest Resource Information Techniques, Chinese Academy of Forestry, Beijing 100091, China.

2 Key Laboratory of Forest Management and Growth Modeling, National Forestry and Grassland Administration, Beijing 100091, China. 3 Experimental Center of Tropical Forestry, Chinese Academy of Forestry, Pingxiang 532600, Guangxi, China. 4 Industry Development and Planning Institute, NFGA, Beijing 100013, China)

Abstract: **【Objective】** Forest carbon and nitrogen storage significantly affect forest ecosystems and global carbon and nitrogen cycles. Forest management can achieve sustainable development by regulating stand structure. Therefore, the relationship between stand structure and soil carbon (SOC) and nitrogen storage (SON) needs in depth study. **【Method】** In this study, experiments were carried out in mixed and pure *Pinus massoniana* forests to analyze the effects of stand density, stand age, and their interaction on the change trends in SOC and SON in different soil layers. **【Result】** The results showed that, in upper (0–20 cm), middle (20–40 cm), and lower (40–60 cm) soil layers, with increased stand density, the SOC of pure *P. massoniana* stands first increased and then decreased, while SON increased monotonically; in mixed *P. massoniana* stands, SOC and SON both increased monotonically. In different development stages (young, middle-aged, and near-mature), the average SOC of pure *P. massoniana* stands were 91.31 t/ha, 88.56 t/ha, and 85.98 t/ha, respectively, while the average SOC of mixed *P. massoniana* stands were 55.92 t/ha, 48.61 t/ha and 55.05 t/ha. The SOC of pure *P. massoniana* stands was significantly higher than mixed *P. massoniana* stands at all growth and development stages. In pure *P. massoniana* stands, with increasing stand density, the SOC of young, middle-aged, and near-mature stands first increased and then decreased, while the SON increased monotonically. In the mixed *P. massoniana* stands, with increasing stand density, the SOC of young, middle-aged, and near-mature stands increased monotonically, while the SON of young stands increased initially and then decreased, while those of middle-aged and near-mature stands increased monotonically. **【Conclusion】** These results emphasized that the artificial regulation of stand density at the appropriate development stage can maximize the carbon and nitrogen fixation potential of forest soil.

Keywords: *Pinus massoniana*; stand structure; stand density; stand age; pure forest; mixed forest; soil carbon storage; soil nitrogen storage.

1 Introduction

The soil carbon and nitrogen pools of forest ecosystems are massive reservoirs that represent more than 70% of global carbon and nitrogen storage (Liu et al., 2007; Pan et al., 2011). Controlling the dynamic changes of soil carbon and nitrogen reserves would be indispensable for alleviating global warming and the sustainable

Funding: This research was funded by the Fundamental Research Funds of CAF(CAFYBB2019ZB005) and the China National Key R&D Program during the 13th Five-year Plan Period(2016YFD060020501).

Correspondence: Xianzhao Liu, lxz9179@163.com

development of forest ecosystems (Li et al., 2013). Forest soil carbon is preserved in many forms, mainly through soil respiration and carbon and nitrogen cycle exchange with aboveground stand and atmosphere. Therefore, a change in forest status will alter forest soil carbon and nitrogen storage (SOC and SON) (Xu and Shang, 2016). The relationship between forest SOC and SON and stand structural characteristics must be understood to effectively manage forest ecosystems. Forest management can be an important driving force in regulating forest ecosystems (Hanan et al., 2018). Indeed, through the proper regulation of stand structure factors, such as density, tree species composition, and age structure, the soil carbon and nitrogen sequestration processes can be controlled to a certain extent (Nordmeyer et al., 2007; Vesterdal et al., 2013; Truax et al., 2018).

In recent years, studies have shown that stand structure has a significant effect on SOC and SON. Stand density is the main stand factor that forest management regulate stands because it affects soil temperature, humidity, and soil respiration rate (Cheng et al., 2015; Lei et al., 2018; Zhang et al., 2018). Stand density is often adjusted via thinning, which decreases stand canopy density, leading to increased stand surface runoff, aggravated soil erosion, and obvious changes in SOC and SON (Jurgensen et al., 2012; Dang et al., 2018). Thinning can also alter the local plant community, so forest management often implements artificial replanting, which forms mixed forests (Ming et al., 2018). Artificial replanting and natural regeneration will directly alter the composition of tree species, making a managed forest significantly different from a natural forest. Tree species composition is another important index of stand structure that affects the content of SOC and SON (Wang et al., 2010). In some cases, rapid growth by individual trees in pure forests can compensate for the initial SOC and SON losses due to thinning (Augusto et al., 2014). Studies have shown that mixed forest diversity is increased by both naturally regenerated trees and artificially planted trees, which can maintain the exchange balance between underground soil carbon and nitrogen and the aboveground stand (Hulvey et al., 2013; Jucker et al., 2014). In addition, mixed forests with diverse compositions can have enriched litter composition and enhanced litter decomposition rates, both of which affect the soil carbon cycle (Sun et al., 2016; Li et al., 2020).

The contents of SOC and SON can be significantly different in different stages of forest growth and development (Wu et al., 2020). In younger stages, the consumption of water and nutrients for tree growth is low, as is the intake of SOC and SON from the soil. With increasing age, the consumption of nutrients by trees increases, and the soil carbon and nitrogen from decomposed litter are also utilized more. The SOC and SON of stands also change dynamically with increasing stand age. In addition, this dynamic may be closely related to changes in stand structure, and may be affected by factors such as tree species composition and stand density (Xia et al., 2021; Na et al., 2021). With the growth of trees, the expansion of tree roots in soil increases correspondingly. With increased soil depth, the carbon and nitrogen in the soil exist in more stable forms that are not easily used by trees and are less affected by trees. Therefore, the influence of stand structure on SOC and SON is different at different depths (Sun et al., 2019). Indeed, topsoil has been seen to be more sensitive to changes in stand structure. The carbon and nitrogen input from litter decomposition and forest root precipitation complete the biochemical process in the surface soil. As it is processed, small amounts of carbon and nitrogen are transported into the deep soil in various forms. The stored carbon and nitrogen in the deep soil mainly comes from the soil itself (Yu et al., 2014; Witing et

al., 2019). However, some studies have found that stand structure does not have different effects on deep soil and surface soil, and the specific respiration rate of deep soil can store unstable carbon and nitrogen in a progressive process, which is a key pathway by which carbon and nitrogen are input into deep soil (Guenet et al., 2012; Heitkotter et al., 2017). Based on the above research, this study explored the relationship between SOCD, SOND, and stand structure in the upper (0–20 cm), middle (20–40 cm), and lower (40–60 cm) soil layers.

Pinus massoniana is the main tree species used for afforestation in barren mountainous areas in subtropical areas. Forest management can improve the *P. massoniana* stand quality by adjusting stand structure (Zhang et al., 2020). Changes in stand structure will affect the exchange between soil and aboveground forest biomass, affect soil respiration rates, and subsequently affect SOCD and SOND. The relationship between the dynamic changes in forest SOCD and SOND and stand structure is of fundamental importance to forest management. Therefore, this study explored the relationship between forest SOCD and SOND and stand structure, primarily to guide and optimize the effectiveness of human Intervention and forest management practices to achieve sustainable forest development.

2 Materials and methods

2.1 Study site

The study was conducted in the Tropical Forestry Experimental Center of Chinese Academy of Forestry, Pingxiang City, Guangxi Province, within latitudes 21°57'47"–22°19'27" N and longitudes 106°39'50"–106°59'30" E. The study area has a subtropical semi-humid-humid climate with distinct dry and wet seasons and abundant light, water, and heat resources. Historically, the mean annual rainfall for the area ranges between 1200–1500 mm and the annual evaporation ranges between 1261–1388 mm. The geomorphology is mainly mountainous and hilly, with an altitude range of 130–1045.9 m, but most areas fall between 500–800 m. The soil-forming parent rocks are mainly argillaceous sandstone, gravelly limestone, granite, and limestone, and the soils are mainly red soil and latosol.

2.2 Experimental design

In order to study the relationship between stand structure and SOCD and SOND, selection of *P. massoniana* evenaged plantation of Tropical Forestry Experimental Center, initial density for 2500 trees/ha. The *P. massoniana* plantation had been managed completely according to the rotation period management mode. We selected 90 standard sample *P. massoniana* plantation plots, including 45 pure *P. massoniana* plots and 45 mixed *P. massoniana* plots. The sample plots were circular plots of 400 m². The mixed forest broad-leaved tree species mainly include *Eucalyptus*, *Mytilaria laosensis*, *Castanopsis hystrix*, etc. *P. massoniana* pure forest sample plots are those tree sample plots with more than 65% of the total stock of *P. massoniana*, and *P. massoniana* mixed forest sample plots are those tree sample plots with 35%–65% of the total stock of *P. massoniana* (Meng, 2006). According to the established requirements of the experiment, the plots were selected so that there was no significant difference between stand factors (i.e., DBH and tree height) and site factors (i.e., altitude, slope, etc.) (Table 1). According to the growth characteristics of *P. massoniana*, the pure and mixed plots could be divided into different age classes: young forest (Mean stand age is 1–10 years), middle-aged forest (Mean stand age is 11–20 years), and near-mature forest (Mean stand age is more than 21 years), with 15 plots in each age group (Meng, 2006).

Table 1. Summary of the characteristics of the sample plots.

Stand type	Proportion of <i>P. massoniana</i>	Initial planting density (trees/ha)	Age group	Mean stand density (trees/ha)	Mean DBH (cm)	Mean H (m)	pH	Elevation (m)	Slope (°)
<i>P. massoniana</i> pure stand	92%	250	Young stand	650	14.83±2.21aA	9.43±1.32aA	4.52±0.35aA	305.95±59.72aA	24.18±3.87aA
			Middle-aged stand	550	18.85±2.31bB	11.73±1.41bB	4.49±0.21aA	316.55±67.23aA	24.12±4.23aA
			Near-mature stand	475	22.73±1.43cC	15.23±1.81cC	4.42±0.28aA	321.48±53.81aA	23.82±2.88aA
<i>P. massoniana</i> mixed stand	49%	250	Young stand	650	14.52±0.67aA	10.43±0.77aA	4.56±0.36aA	305.45±63.75aA	24.59±4.53aA
			Middle-aged stand	575	17.85±1.22bB	11.23±1.09aB	4.63±0.28aA	314.22±61.34aA	23.73±4.26aA
			Near-mature stand	500	21.29±2.34cC	14.95±2.63cC	4.51±0.26aA	324.17±43.74aA	24.62±3.52aA

Note: different lowercase letters "a, b, c" indicate that there are significant differences, and different capital letters "A, B, C" indicate significant differences in *P. massoniana* pure stand and *P. massoniana* mixed stand.

2.3 Field survey and soil sampling

In sample plots where trees have $DBH \geq 5$ cm and young trees have heights > 30 cm and $DBH < 5$ cm, the DBH and tree height can be measured using the DBH ruler and ultrasonic tree altimeter. For each plot, the tree species names were recorded, the DBH and tree heights were measured, and the shrub species, numbers, and heights were recorded. Soil samples were collected according to the plum blossom method, and 1 kg of the upper, middle, and lower soils were collected according to 20 cm intervals in the east, west, south, north, and center of each plot area. The upper layer of soil was 0–20 cm, the middle layer of soil was 20–40 cm, and the lower layer of soil was 40–60 cm. The samples were air-dried and crushed, and then the fine roots of the plants were removed and samples were sealed for storage. Soil organic carbon was determined using the potassium dichromate method; and available nitrogen was determined via alkaline hydrolysis distillation (Qiu et al., 2019).

2.4 Data processing

We calculated the SOCD as follows (Kong et al., 2019):

$$SOCD = SOC \times h \times \rho \times 10 \quad (1)$$

$$SOND = N_1 \times h \times \rho \times 10 \quad (2)$$

where SOCD is soil organic carbon storage (t/ha); SOND is soil nitrogen storage (t/ha); SOC is soil organic carbon content (g/kg); N_1 is soil organic nitrogen content (g/kg); h is soil thickness ($h = 0.2$ m); and ρ is soil bulk density (g/cm^3).

First, a multi-factor variance analysis was conducted to analyze whether stand density, age group, and soil layer had significant effects on SOCD and SOND. Second, a linear mixed model was adopted to consider the effects of age group, soil layer, and their interaction on SOCD and SOND. The expressions were as follows (Hernández et al.,

2016):

$$\begin{aligned}
 Y_{ij} &= a_0 + \mu_j + a_1 \times N_{ij} + \varepsilon_{ij} \\
 Y_{ik} &= a_0 + \mu_k + a_1 \times N_{ik} + \varepsilon_{ik} \\
 Y_{ijk} &= a_0 + \mu_j \times \mu_k + a_1 \times N_{ijk} + \varepsilon_{ijk}
 \end{aligned}
 \tag{3}$$

where a_0 and a_1 are the fixed parameters of the model; μ_j and μ_k are the random effects; ε_{ij} , ε_{ik} and ε_{ijk} are error term; Y_{ij} is the logarithmic conversion value of SOCD or SONND at i^{th} sample of the j^{th} age group; Y_{ik} is the logarithmic conversion value of SOCD or SONND at i^{th} sample of the k^{th} soil layer; and Y_{ijk} is the logarithmic conversion value of SOCD or SONND at i^{th} sample of the k^{th} soil layer and the j^{th} age group. N_{ij} is the stand density at i^{th} sample of the j^{th} age group; N_{ik} is the stand density at i^{th} sample of the k^{th} soil layer; N_{ijk} is the stand density at i^{th} sample point of the k^{th} soil layer and the j^{th} age group. Age group, soil layer, and stand density are nested structure, such as in the same age group, several soil layers are nested in the same age group; in the same age group and soil layer, several stand density levels are nested in the same age group and soil layer.

Since the defined domain of the independent variable (stand density) in this study was small, a logarithmic conversion of soil carbon and nitrogen reserves was carried out, and the logarithmic conversion values of the soil carbon and nitrogen reserves were used to eliminate any autocorrelation or heteroscedasticity in the data. Data analysis and illustrations were completed in R 4.0.3 and the OriginPro software, respectively.

3 Results

3.1 Factors affecting SOCD and SONND

The multivariate analysis of variance (Table 2) showed that stand density and soil depth had significant effects on SOCD and SONND ($P < 0.001$). The interactions between stand age and soil depth and between stand density and soil depth significantly affected SOCD and SONND. Stand density and soil depth both significantly affected SOCD and SONND, the former above ground and the later underground, and the interaction between them was more sensitive than the interaction between stand age and soil depth.

Table 2. Analysis results affecting SOCD and N. In the table, asterisk represents statistical significance.

Factor	SOCD		N	
	F Value	P Value	F Value	P Value
Stand age	0.0472	0.8284	1.5511	0.2148
Stand density	19.7672	<0.001***	7.6151	0.0065**
Soil depth	79.3354	<0.001***	57.9299	<0.001***
Stand age×Stand density	0.1232	0.7261	2.9043	0.0903
Stand age×Soil depth	1.2457	0.0345*	3.5624	0.0475*
Stand density×Soil depth	1.5763	0.0241*	4.1352	0.0286*

* $P < 0.05$, ** $P < 0.01$, and *** $P < 0.001$.

3.2 SOCD and SONND at different depths in mixed and pure *P. massoniana* stands of different densities

The linear regression was able to accurately describe the effect of stand density on SOCD at different depths (Figure 1a, 1b). The change trends of SOCD in the upper, middle, and lower layers were similar to the trend with decreasing stand density. In the pure *P. massoniana* stands, with increased stand density, the SOCD in the upper,

middle, and lower layers increased at first and then decreased. When the stand density was less than 1500 trees/ha, the growth rate of SOCD in the upper layer was significantly higher than those in the middle and lower layers. When the stand density was 1500–2000 trees/ha, upper layer, the SOCD in lower layer and middle layer reached the peak respectively. When stand density exceeded 1500 trees/ha, the SOCD in upper, middle, and lower layers decreased with increasing stand density, and the rate of SOCD decrease in the upper layer was the fastest. In mixed *P. massoniana* stands, the SOCD of the upper, middle, and lower layers increased uniformly with increasing stand density. When the stand density increased to 2500 trees/ha, the increases in SOCD in the upper, middle, and lower layers were 94.07 t/ha, 53.07 t/ha, and 29.83 t/ha, respectively.




Figure 1. SOCD at different depths and stand densities. (a) SOCD at different depths and different stand densities in pure *P. massoniana* stands; (b) SOCD at different depths and different stand densities in mixed *P. massoniana* stand.

The SOND of upper, middle, and lower layers showed a “J” curve, and the SOND increased with increasing stand density (Figure 2a, 2b). The N contents of upper and middle layers were higher in pure *P. massoniana* stands than in mixed *P. massoniana* stands, while the SOND content of the lower layer was lower in pure *P. massoniana* stands than in mixed *P. massoniana* stands. With increasing stand density, the difference in the SOND contents of the upper and middle layers of the pure and mixed *P. massoniana* stands became larger, while the SOND content in the lower layer of the pure *P. massoniana* stand gradually increased and eventually surpassed that in the mixed stand. When the stand density increased to 2500 trees/ha, the SOND contents in the upper, middle, and lower layers of the pure *P. massoniana* stand increased by 2.86 g/kg, 2.15 g/kg, and 1.33 g/kg, respectively, while those of the mixed *P. massoniana* stands increased by 1.63 g/kg, 1.13 g/kg, and 0.91 g/kg.




Figure 2. SOND at different depths and stand densities (a) SOND at different depths and stand densities in pure *P. massoniana* stands; (b) SOND at different depths and different stand densities in mixed *P. massoniana* stands.

3.3 Effects of age group on SOCD and SOND at different depths in pure and mixed stands of *P. massoniana*

According to the analysis of variance (Table 3; Figure 5a, 5b, 5c, 5d), there were significant differences in SOCD among the different soil layers in different age groups of pure and mixed *P. massoniana* stands. The SOCD of the upper, middle, and lower layers of the pure *P. massoniana* stands were significantly higher than those of the mixed *P. massoniana* stands. The SOND contents in the upper layers of both pure and mixed *P. massoniana* stands were significantly different from those of the middle and lower layers, and there were no significant differences between middle and lower layers. There were no significant differences in SOND content between the upper, middle, and lower layers of either pure or mixed *P. massoniana* stands.

Table 3. Analysis results of SOCD and N of different age groups and different depths in pure and mixed stands of *P. massoniana*.

Stand type	Soil depth (cm)	SOCD(t/ha)			N(t/ha)		
		Young stand	Middle-aged stand	near-mature stand	Young stand	Middle-aged stand	near-mature stand
<i>P. massoniana</i> pure stand	A	137.74±14.57a A	148.13±17.51aA	141.61±26.06a A	1.95±0.48a A	1.56±0.64a A	1.81±0.39a A
	AB	82.52±6.34bA	71.43±13.61bA	70.19±11.51bA	1.11±0.57b A	1.13±0.31b A	1.16±0.44b A
	B	53.64±5.76cA	46.12±2.75cA	46.14±8.74cA	0.97±0.65b A	0.95±0.29b A	0.81±0.17b A
<i>P. massoniana</i> mixed stand	A	87.51±4.24aB	79.36±8.52aB	89.02±3.48aB	1.53±0.51a A	1.45±0.28a A	1.49±0.24a A
	AB	51.91±3.49bB	42.23±6.02bB	47.95±5.77bB	1.11±0.45b A	0.99±0.21b A	0.91±0.21b A
	B	28.35±2.61cB	24.23±4.83cB	28.17±1.56cB	0.87±0.32b A	0.84±0.22b A	0.79±0.24b A

Note: values are the mean ± standard deviation (SD), different lowercase letters "a, b, c" indicate that there are significant differences in SOCD and N reserves in different soil layers, and different capital letters "A, B, C" indicate significant differences in SOCD and N reserves in different stand types.

The SOCD decreased with increasing soil depth in all different age groups of both pure and mixed *P. massoniana* stands, albeit at different rates (Figure 3a, 3b, 3c). The changes in SOCD of different age groups in the upper, middle, and lower layers were consistent with the trends observed with changes in stand density. When the stand density was 1000–1500 trees/ha, the SOCD of the middle-aged *P. massoniana* pure stand peaked. At stand densities of 1500–2000 trees/ha and 2000–2500 trees/ha, the SOCD of the near-mature and young stands peaked, respectively. Compared with the initial SOCD, the SOCD peaks of the middle-aged, young, and near-mature pure *P. massoniana* stands increased by 3.5, 1.7, and 4.1 times, respectively. When the stand density increased to 2500 trees/ha, the SOCD of middle-aged, young, and near-mature mixed *P. massoniana* stands increased by 0.85, 1.59, and 2.69 times, respectively.




Figure 3. SOCD with different stand densities and different age groups at different depths in pure and mixed *P. massoniana* stands. (a) SOCD in the upper soil layer with different stand densities and different age groups in mixed and pure *P. massoniana* stands; (b) SOCD in the middle soil layer with different stand densities and different age groups in mixed and pure *P. massoniana* stands; (c) SOCD in the lower soil layer with different stand densities and different age groups in mixed and pure *P. massoniana* stands.

The SOND contents of different age groups of pure and mixed *P. massoniana* stands decreased with increasing soil depth (Figure 4a, 4b, 4c). Overall, in the upper, middle, and lower layers, the SOND contents of different age groups of the pure *P. massoniana* stands increased with increasing stand density. The young mixed *P. massoniana* stands increased initially, but then decreased with increasing stand density, while the middle-aged and near-mature stands increased continuously. With increasing stand density, the SOND contents of young pure *P. massoniana* stands were always higher than those of the mixed *P. massoniana* stands in the upper and middle layers, but at densities between 500–1200 trees/ha, the SOND content of the middle layers of mixed *P. massoniana* stands were briefly higher than the middle layer of the young pure *P. massoniana* stands.




Figure 4. SOND with different stand densities and different age groups at different depths in pure and mixed *P. massoniana* stands. (a) SOND in the upper soil layer with different stand densities and different age groups in mixed and pure *P. massoniana* stands; (b) SOND in the middle soil layer with different stand densities and different age groups in mixed and pure *P. massoniana* stands; (c) SOND in the lower soil layer with different stand densities and different age groups in mixed and pure *P. massoniana* stands.




Figure 5. Analysis of variance results. (a) 95% family-wise confidence level for SOCD in different age groups in different soil layers; (b) 95% family-wise confidence level of SOND in different age groups; (c) 95% family-wise confidence level of SOCD in mixed and pure *P. massoniana* stands; (d) 95% family-wise confidence level of SOND in mixed and pure *P. massoniana* stands.

4 Discussion

4.1 Effect of stand density on SOCD and SOND of pure and mixed *P. massoniana* stands

Forest type, stand density, and age group structure do not provide a sufficiently comprehensive assessment for effective forest management, so other important influencing factors like the relationships between soil carbon and nitrogen and stand structure must be considered (Ali et al., 2019). The SOCD of pure *P. massoniana* stands and SOND of mixed *P. massoniana* stands of different densities exhibited significant differences. The change trends of SOCD in pure and mixed *P. massoniana* stands with different densities were different, while the change trends of SOND were roughly similar across different stand densities. With increases in stand density, the SOCD of pure *P. massoniana* stands initially increased, but then decreased, while the SOCD of mixed *P. massoniana* stands increased monotonically. As for the interpretation rate of biological characteristics, the interpretation rate of SOCD was higher and tended to be more stable at higher stand densities. Most research to date has shown that stand density has a significant effect on SOCD, but the effect of stand density on SOCD is indirect, acting through its direct effect on soil respiration (Kuzyakov and Gavrichkova, 2010; Olajuyigbe et al., 2012; Bujalský et al., 2014). Soil respiration is an important soil carbon loss pathway and it is significantly affected by climate characteristics such as soil temperature and humidity. The regulation of stand density will produce obvious changes in local microclimates, therefore soil respiration will be significantly enhanced or weakened. When the stand density is too high or too low, soil respiration can be restricted. Therefore, through soil respiration, stand density will indirectly affect soil carbon storage. As stand density increases, soil respiration will increase until peaking, at which point the soil carbon storage will also peak; continued increases in stand density will then lead to decreases in respiration and soil carbon storage,

resulting in an “S” curve change trend (Colman and Schimel, 2013; Shao et al., 2013; Liu et al., 2021). The difference of soil aggregation size distribution indicates that plant root input and litter decomposition play a key role in the change of SOCD (Oades, 1993). Macroaggregates can absorb carbon quickly, while microaggregates absorb carbon slowly, but the storage form of carbon will be more stable (Li et al., 2013). The pure forest has a single tree species, and most of the soil is dominated by macroaggregates, while mixed forest has larger soil diversity and soil pore space, which makes it easier for soil to form micro poly collective. With the increase of stand density, the macroaggregates in pure *P. massoniana* forest soil quickly absorb the carbon decomposed by litter, which leads to the increase of SOCD. However, coniferous species mainly produce low-quality litter, and there are many substances that are difficult to decompose (such as lignin, cellulose, etc.) (Shi et al., 2018). A small amount of micro poly collectively absorb carbon more slowly, and the soil carbon absorbed by macro-aggregates is unstable and gradually loses, which leads to the decrease of SOCD. In the mixed forest, fast-growing broad-leaved tree species such as *Eucalyptus*, *Mytilaria laosensis*, *Castanopsis hystrix*, etc. mainly produce high-quality litter. The litter accumulated by the increase of stand density is slowly decomposed and absorbed by a large number of microaggregates, forming stable soil carbon (Contrufo et al., 2019; Lavalley et al., 2020), and the SOCD increases monotonically.

The SOCD and SOND of mixed *P. massoniana* stands increased with increasing stand density, which was consistent with the study by Laganier (2017) of the soil carbon cycle balance, which observed a specific “threshold” at which capacity limitations were reached at the input and output ends of the carbon and nitrogen cycles. According to the afforestation principle of mixed forests, with increased or decreased stand density, the complementary tree species in mixed stands will show relative increases or decreases, which is the theoretical equivalent to enlarging or reducing the specific “threshold” of soil carbon and nitrogen cycles according to a specific multiplier, while the SOCD and SOND showed relative exponential growth, or “J”, curves (Son et al., 1999; Kuzyakov and Larionova, 2005). With increasing stand density, mixed *P. massoniana* stands became predominantly composed of small-diameter trees, which accounted for ~56%. In the mixed *P. massoniana* stands, most of the naturally regenerated tree species that met the standards of imported wood were tree species considered complementary for afforestation. As the undergrowth seedlings increasingly meet the standards for imported wood, the carbon fixation and nitrogen fixation in forest ecosystem will increase, and the SOCD and SOND reserves will also naturally increase. Mixed *P. massoniana* stands have a stronger resilience to climate change disturbance because of their tree species diversity and stable stand structures. Local climate changes caused by the regulation of stand density can promote the cyclic conversion between understory vegetation and soil carbon, thus contributing to the long-term preservation of SOCD (Silva et al., 2015).

4.2 Differences in SOCD and SOND between pure and mixed *P. massoniana* stands

The SOCD of the pure *P. massoniana* stands was significantly higher than that of the mixed *P. massoniana* stands, while the difference in SOND was not significant. This conclusion was different from previously published reports (e.g., Guckland et al., 2009; Schleuß et al., 2014). Specifically, comparing pure to mixed *P. massoniana* stands, it could be seen that artificially replanted and naturally renewed broad-leaved tree species grew slower than

P. massoniana, which is a fast-growing tree species. Therefore, the pure stands could form a main forest layer more quickly, thereby reducing light penetration to the understory and inhibiting the growth of herbs and shrubs, which is matched by corresponding increases in litter and SOCD (Dawud et al., 2017; Terrer et al., 2021). However, as broad-leaved species of mixed stand gradually grow into the main forest layer, the canopy density gradually increases and exceeds that of pure stands, and the SOCD of the mixed stand will also increase. In this study, while the soil in the upper layer was strongly affected by surface runoff, the differences in the SOCD of the pure and mixed stands in the middle and lower layers decreased with increasing age, from 30.61 to 22.24 and from 25.29 to 17.97. This conclusion was supported by previous studies which have shown that the characteristics of species or existing functional groups significantly impact soil carbon storage, with coniferous species having a stronger adaptability to environmental change than broad-leaved species, and coniferous species also having a more positive impact on root biomass (soil carbon input), which can promote increases in SOCD (Dawud et al., 2016; Ratcliffe et al., 2017; Finér et al., 2017).

4.3 Effect of stand age on SOCD and SOND in pure and mixed stands of *P. massoniana*

In this paper, the differences in SOCD and SOND between pure and mixed stands of *P. massoniana* in different age groups and their change trends with changing stand density were studied. The SOCD of different age groups of *P. massoniana* pure stand increased initially and then decreased with increasing stand density, and the SOCD of different age groups of mixed *P. massoniana* stands increased with increasing stand density. Among the peaks of SOCD of each age group of the pure *P. massoniana* stands, that of the near-mature stand was highest, possibly because there was obvious difference in the number of large-diameter trees among different age groups. However, with increasing age, the litter layer will form a thicker layer of coarse humus and the abundance of strong enzymatically-active microorganisms will be higher, which will enhance the soil carbon cycle and increase the soil carbon content (Prieto et al., 2012; Stevenson et al., 2017; Xu et al., 2021). The N of young mixed *P. massoniana* stands increased initially and then decreased with increasing stand density. In the young stands the canopy was in the developing stage, this meant that the canopy density was low and the shading effect in the upper story was poor, so the water and light resources in the young stands were relatively abundant. The SOND content increased gradually as soil ammoniation and nitrification were promoted. With increasing stand density, tree crown growth, canopy density, and upper shading effect gradually increased, but soil ammoniation and nitrification slowed as resources became limited, thus N decreased. Deep soil nitrogen exists in a more stable form, and its sensitivity to aboveground stand changes gradually decreases with depth (Mayer et al., 2020).

5 Conclusions

Through qualitative analysis of linear mixed models, it was found that stand density had a significant impact on SOCD and SOND in pure and mixed stands of *P. massoniana*. With increasing stand density, SOCD and SOND in pure *P. massoniana* stands first increased and then decreased, while SOCD and SOND in mixed *P. massoniana* stands increased monotonically. During the growth and development of trees, the SOCD and SOND trends in young, middle-aged, and near-mature pure and mixed *P. massoniana* stands were consistent with the increases in stand density. For SOND content, that of young mixed *P. massoniana* stands first increased and then decreased, while

those of middle-aged and near-mature stands increased monotonically. Taking corresponding management measures in the appropriate growth and development stages and regulating stand density can therefore effectively realize the maximum carbon and nitrogen fixation potential of the stand, thus maximizing the SOCD and SOND of the stand. There were significant differences in SOCD of different stand types. This paper showed that the SOCD of pure *P. massoniana* stands in each growth and development stage was significantly higher than that of mixed stands. Under the rotation management scheme, pure *P. massoniana* stand has higher storage potential, which further shows that regulating stand density can directly and effectively affect SOCD.

References

- Ali, A, Dong, D, Akhtar, K, Teng, M, et al. 2019. Response of understory vegetation, tree regeneration, and soil quality to manipulated stand density in a *Pinus massoniana* plantation. *Global Ecology and Conservation*, 20, e00775, doi: 10.1016/j.gecco.2019.e00775.
- Augusto, L, Davies, T, Delzon, S, et al. 2014. The enigma of the rise of angiosperms: can we untie the knot? *Ecology Letters*, 17, 1326-1338, doi: 10.1111/ele.12323.
- Bujalský, L, Kaneda, S, Dvorsck, P, Frouz, J. 2014. In situ soil respiration at reclaimed and unreclaimed post-mining sites: Responses to temperature and reclamation treatment. *Ecological Engineering*. 68, 53-59.
- Chen, S.T, Zou, J.W, Hu, Z.H, et al. 2014. Global annual soil respiration in relation to climate, soil properties and vegetation characteristics: Summary of available data. *Agricultural & Forest Meteorology*. 198, 335-346.
- Cheng, X.Q, Kang F.F, Han H.R, et al. 2015. Effect of thinning on partitioned soil respiration in a young *Pinus tabulaeformis* plantation during growing season. *Agricultural and Forest Meteorology*. 214-215, 473-482.
- Colman, B.P, Schimel, J.P. 2013. Drivers of microbial respiration and net N mineralization at the continental scale. *Soil Biology & Biochemistry*. 60, 65-76.
- Cotrufo, M.F, Ranalli, M.G, Haddix, M.L, et al. 2019. Soil carbon storage informed by particulate and mineral-associated organic matter. *Nature Geoscience*. 12, 989-994.
- Dang, P, Gao, Y, Liu, J.L, et al. 2018. Effects of thinning intensity on understory vegetation and soil microbial communities of a mature Chinese pine plantation in the Loess Plateau. *Science of the Total Environment*. 630.
- Dawud, S.M, Raulund-Rasmussen, K, Domisch, T, et al. 2016. Is tree species diversity or species identity the more important driver of soil carbon stocks, C/N ratio, and pH? *Ecosystems*. 19, 645-660.
- Dawud, S.M, Raulund-Rasmussen, K, Ratcliffe, S, et al. 2017. Tree species functional group is a more important driver of soil properties than tree species diversity across major European forest types. *Functional Ecology*. 31, 1153-1162.
- Finér, L, Domisch, T, Dawud, S.M, et al. 2017. Conifer proportion explains fine root biomass more than tree species diversity and site factors in major European forest types. *Forest Ecology & Management*. 406, 330-350.
- Guckland, A, Jacob, M, Flessa, H, et al. 2009. Acidity, nutrient stocks, and organic-matter content in soils of a temperate deciduous forest with different abundance of European beech (*Fagus sylvatica* L.). *Journal of Plant Nutrition and Soil Science*. 172, 500-511.
- Guenet, B, Juarez, S, Bardoux, G, et al. 2012. Evidence that stable C is as vulnerable to priming effect as is more labile C in soil. *Soil Biology and Biochemistry*. 52, 43-48.
- Hanan, E.J, Tague, C, Choate, J, et al. 2018. Accounting for disturbance history in models :using remote sensing to constrain carbon and nitrogen pool spin-up. *Ecological Applications*. 28, 1197-1214.
- Heitkotter, J, Niebuhr, J, Heinze, S, et al. 2017. Patterns of nitrogen and citric acid induced changes in C-turnover and enzyme activities are different in topsoil and subsoils of a sandy Cambisol. *Geoderma*. 292, 111-117.
- Hernández, J, Pino, A, Vance, E.D, et al. 2016. Eucalyptus and *Pinus* stand density effects on soil carbon sequestration. *Forest Ecology and Management*. 368, 28-38.
- Hulvey, K.B, Hobbs, R.J, Standish, R.J, et al. 2013. Benefits of tree mixes in carbon plantings. *Nature Climate Change*. 3, 869-874.
- Jucker, T, Bouriaud, O, Avacaritei, D, et al. 2014. Stabilizing effects of diversity on aboveground wood production in forest ecosystems: linking patterns and processes. *Ecology Letters*. 17, 1560-1569.
- Jurgensen, M, Tarpey, R, Picjens, J, et al. 2012. Long-term Effect of Silvicultural Thinnings on Soil Carbon and Nitrogen Pools. *Soil Science Society of America Journal*. 76, 1418-1425.

- Kong, J.Q, Du, Z.Y, Yang, R, et al. 2019. Evolutionary characteristics of soil organic carbon storage in soil plough layer under a crop- land reclamation process in desert oasis. *Chinese Journal of Applied Ecology*. 30, 180-188.(In Chinese)
- Kuzyakov, Y, Gavrichkova, O. 2010. REVIEW: Time lag between photosynthesis and carbon dioxide efflux from soil: a review of mechanisms and controls Y. KUZYSKOV & O. GAVRICHKOVA TIME LAG BETWEEN PHOTOSYNTHESIS AND CO EFFLUX FROM SOIL. *Global Change Biology*. 16, 3386-3406.
- Kuzyakov, Y, Larionova, A.A. 2005. Root and rhizomicrobial respiration: a review of approaches to estimate respiration by autotrophic and heterotrophic organisms in soil. *Journal of Plant Nutrition and Soil Science*. 168, 503-520.
- Lavallee, J.M, Soong, J.L, Cotrufo, M.F. 2020. Conceptualizing soil organic matter into particulate and mineral-associated forms to address global change in the 21st century. *Global Change Biology*. 26, 261-273.
- Laganiere, J, Pare, D, Thiffault, E, Bernier, P.Y. 2017. Range and uncertainties in estimating delays in greenhouse gas mitigation potential of forest bioenergy sourced from Canadian forests. *GCB Bioenergy*. 9, 358-369.
- Lei, L, Xiao, W.F, Zeng, L.X, et al. 2018. Thinning but not understory removal increased heterotrophic respiration and total soil respiration in *Pinus massoniana* stands. *The Science of the total environment*. 621, 1360-1369.
- Li, M.M, Zhang, X.C, Pang, G. et al. 2013. The estimation of soil organic carbon distribution and storage in a small catchment area of the Loess Plateau. *Catena*. 101, 11-16.
- Li, Y, Liu, X.J, Xu, W.B, et al. 2020. Effects of diversity, climate and litter on soil organic carbon storage in subtropical forests. *Forest Ecology & Management*. 476, 118479.
- Li, Y.L, Yang, F.F, Ou, Y.X, et al. 2013. Changes in forest soil properties in different successional stages in lower tropical China. *Plos One*. 8, 11, e81359.
- Liu, M.Q, Hu, F, Chen, X.Y. 2007. A review on mechanisms of soil organic carbon stabilization. *Acta Ecologica Sinica*. 27, 2642-2650.(In Chinese)
- Liu, T.R, Peng, D.L, Tan, Z.J, et al. 2021. Effects of stand density on soil respiration and labile organic carbon in different aged *Larix principis-rupprechtii* plantations. *Ecological Processes*. 10, 1-15.
- Mayer, M, Prescott, C.E, Abaker, W.E.A, et al. 2020. Tamm Review: Influence of forest management activities on soil organic carbon stocks: A knowledge synthesis. *Forest Ecology and Management*, 466, 118127.
- Meng, X.Y. *Forest Mensuration*, 3rd ed, China Forestry Publishing House, Beijing, China, 2006. (In Chinese)
- Ming, A.G, Yang, Y.J, Liu, S.R, et al. 2018. Effects of Near Natural Forest Management on Soil Greenhouse Gas Flux in *Pinus massoniana* (Lamb.) and *Cunninghamia lanceolata* (Lamb.) Hook. *Plantations. Forests*. 9, 229.
- Na, M, Sun, X.Y, Zhang, Y.D, et al. 2021. Higher stand densities can promote soil carbon storage after conversion of temperate mixed natural forests to larch plantations. *European Journal of Forest Research*. 140, 373-386.
- Nordmeyer, A, Henley, D, Watt, M, et al. 2007. Ecosystem carbon accretion 10 years after afforestation of depleted subhumid grassland planted with three densities of *Pinus nigra*. *Global Change Biology*. 13, 1414-1422.
- Oades, J.M. 1993. The role of biology in the formation, stabilization and degradation of soil structure - ScienceDirect. *Soil Structure/Soil Biota Interrelationships*. 56, 377-400.
- Olajuyigbe, S, Tobin, B, Saunders, M, et al. 2012. Forest thinning and soil respiration in a Sitka spruce forest in Ireland. *Agricultural and Forest Meteorology*. 157, 86-95.
- Pan, Y.D, Birdsey, R.A, Fang, J.Y, et al. 2011. A large and persistent carbon sink in the world's forests. *Science*. 333, 988-993.
- Prieto, I, Armas, C, Pugnaire, F.I. 2012. Water release through plant roots: new insights into its consequences at the plant and ecosystem level. *The New Phytologist*. 193, 830-841.
- Qiu, X.C, Peng, D.L, Wang, H.B, et al. 2019. Minimum data set for evaluation of stand density effects on soil quality in *Larix principis-rupprechtii* plantations in North China. *Ecological Indicators*. 103, 236-247.
- Ratcliffe, S, Wirth, C, Jucker, T, et al. 2017. Biodiversity and ecosystem functioning relations in European forests depend on environmental context. *Ecology Letters*. 20, 1414-1426.
- Schleuß, P.M, Heitkamp, F, Leuschner, C, et al. 2014. Higher subsoil carbon storage in species-rich than species-poor temperate forests. *Environmental Research Letters*. 9, 014007.
- Shao, P, Zeng, X.B, Moore, D.J.P, et al. 2013. Soil microbial respiration from observations and Earth System Models. *Environmental Research Letters*. 8, 034034.
- Shi, F.J, Huang, Z.Y, Li, T, et al. 2015. Vertical Distribution and correlation Analysis of soil microbial Biomass carbon and nitrogen in Wangtian Tree Natural Forest. *Forestry and Environmental Science*. 2018, 34, 5, doi: 10.3969/j.issn.1006-4427.2018.06.012.(In Chinese)
- Silva, P.M, Rammer, W, Seidl, R, Tree species diversity mitigates disturbance impacts on the forest carbon cycle. *Oecologia*. 177, 619-630.

- Son, Y.H, Lee, W.K, Lee, S.E, et al. 1999. Effects of thinning on soil nitrogen mineralization in a Japanese larch plantation. *Communications in Soil Science and Plant Analysis*. 30, 2539-2550.
- Stevenson, B.A, Hunter, D.W, Rhodes, P.L. 2014. Temporal and seasonal change in microbial community structure of an undisturbed, disturbed, and carbon-amended pasture soil. *Soil Biology & Biochemistry*. 75, 175-185.
- Sun, X.D, Wang, w.n, Razaq, M, Sun, H.L. 2019. Effects of stand density on soil organic carbon storage in the top and deep soil layers of *Fraxinus mandshurica* plantations. *Austrian Journal of Forest Science*. 136, 27-44.
- Sun, X.L, Zhao, J, You, Y.M, et al. Soil microbial responses to forest floor litter manipulation and nitrogen addition in a mixed-wood forest of northern China. *Scientific Reports*. 2016, 6, 19536.
- Terrer, C, Phillips, R.P, Hungate, B.A, et al. 2021. trade-off between plant and soil carbon storage under elevated CO₂. *Nature*. 591, 599-603.
- Truax, B, Fortier, J, Gagnon, D, et al. 2018. Planting density and site effects on stem dimensions, stand productivity, biomass partitioning, carbon stocks and soil nutrient supply in hybrid poplar plantations. *Forests*. 9, 293.
- Vesterdal, L, Clarke, N, Sigurdsson, B.D, et al. 2013. Do tree species influence soil carbon stocks in temperate and boreal forests? *Forest Ecology & Management*. 309, 4-18.
- Wang, H, Huang, Y, Wang, S.L, et al. 2010. Carbon and nitrogen storage under different forest ecosystems in mid-subtropical regions. *Chinese Journal of Eco-Agriculture*. 18, 576-580.(In Chinese)
- Witing, F, Gebel, M, Kurzer, H.J, et al. 2019. Large-scale integrated assessment of soil carbon and organic matter-related nitrogen fluxes in Saxony (Germany). *Journal of Environmental Management*. 237, 272-280.
- Wu, H.L, Xiang, W.H, Ouyang, S, et al. 2020. Tree growth rate and soil nutrient status determine the shift in nutrient-use strategy of Chinese fir plantations along a chronosequence. *Forest Ecology & Management*. 460, 117896.
- Xia, Q, Chen, L, Xiang, W.H, et al. 2021. Increase of soil nitrogen availability and recycling with stand age of Chinese fir plantations. *Forest Ecology and Management*. 480, 118643.
- Xu, C.H, Xu, X, Ju, C.H, et al. 2021. Long-term, amplified responses of soil organic carbon to nitrogen addition worldwide. *Global change biology*. 27, 1170-1180.
- Xu, M, Shang, H. 2016. Contribution of soil respiration to the global carbon equation. *Journal of Plant Physiology*. 203, 16-2.
- Yu, S.B, Wang, D, Dai, W, et al. 2014. Soil carbon budget in different-aged Chinese fir plantations in south China(Article). *Journal of Forestry Research*. 25, 621-626.
- Zhang, K.X, Liu, X.Z, Lei, X.D, et al. 2020. Short-term economic benefit evaluation of different management modes of *Pinus massoniana* plantation. *Journal of Beijing Forestry University*. 44, 43-54. (In Chinese)
- Zhang, X.Z, Guan, D.X, Li, W.B, et al. 2018. The effects of forest thinning on soil carbon stocks and dynamics: A meta_analysis. *Forest Ecology & Management*. 429, 36-43.